

UNIVERSITY OF BAHRAIN
 COLLEGE OF SCIENCE
 PHYSICS DEPARTMENT


PHYCS 102
 TEST 3

DATE: 2/1/2001


TIME: 55 MIN.

NAME:	ID#:	SECTION:
-------	------	----------

- Q1.** In the circuit shown the capacitor is fully charged. Then, at $t = 0$ the switch is thrown from “a” to “b”. This causes the current to decrease to 0.5 of its initial value in $40 \mu\text{s}$.
- Calculate the value of R .
 - What is the value of the capacitor charge Q at $t = 0$?
 - What is the value of Q at $t = 60 \mu\text{s}$?




- Q2.** In the figure shown below, a circular loop of radius $R = 20\text{cm}$ carries a current $I_1 = 2\text{A}$ and a very long straight wire carries a current $I_2 = 5\text{A}$. Use superposition method to determine the magnitude and direction of the total magnetic field at the center “O” of the loop.




- Q3.** A proton is accelerated by 56KV , enters a uniform magnetic field (\vec{B}) in a direction perpendicular to (\vec{B}). The proton moves in a circular path of radius 8m . Determine :
- the magnitude of \vec{B} ,
 - the time required to make 5 revolutions.

- Q4.** A circular loop of radius $R = 10 \text{ cm}$ consists of 50 closely wrapped turns in which each carries a current of 0.5A . The loop is placed in a uniform magnetic field of $B = 0.4\text{T}$ directed in the positive x -axis, as shown in the figure.
- What is the resultant magnetic force on the loop?
 - Calculate the magnitude of the torque $\vec{\tau}$ on the loop.
 - What is the direction of $\vec{\tau}$? Describe the expected rotation of the loop?



- Q5.** A short straight wire of length $L=0.3\text{m}$ carries a current $I_2 = 2\text{A}$ is placed perpendicular at a distance $d= 0.1\text{m}$ near a long straight wire that carries a current $I_1 = 3\text{A}$ as shown in the figure. Determine the magnitude and direction of magnetic force that exerted on the short wire.



$$m_p = 1.67 \times 10^{-27} \text{Kg.}$$

$$E = 1.6 \times 10^{-19} \text{C}$$

Good luck

3. A charged spherical shell of radius R has a total charge Q placed inside an uncharged conducting spherical shell that has an inner radius a and outer radius b . **Find :**
- a) The electric field every where, i.e. in each region 1,2,3 and 4.
 - b) The induced surface charge densities on the inner and outer surfaces of the uncharged conducting spherical shell.

